

Equivariant multiplicities of Coxeter arrangements and invariant bases

Takuro Abe ^{*} Hiroaki Terao [†] Atsushi Wakamiko [‡]

December 30, 2010

Abstract

Let \mathcal{A} be an irreducible Coxeter arrangement and W be its Coxeter group. Then W naturally acts on \mathcal{A} . A multiplicity $\mathbf{m} : \mathcal{A} \rightarrow \mathbb{Z}$ is said to be equivariant when \mathbf{m} is constant on each W -orbit of \mathcal{A} . In this article, we prove that the multi-derivation module $D(\mathcal{A}, \mathbf{m})$ is a free module whenever \mathbf{m} is equivariant by explicitly constructing a basis, which generalizes the main theorem of [T2002]. The main tool is a primitive derivation and its covariant derivative. Moreover, we show that the W -invariant part $D(\mathcal{A}, \mathbf{m})^W$ for any multiplicity \mathbf{m} is a free module over the W -invariant subring.

1 Introduction

Let V be an ℓ -dimensional Euclidean space with an inner product $I : V \times V \rightarrow \mathbb{R}$. Let S denote the symmetric algebra of the dual space V^* and F be its quotient field. Let Der_S be the S -module of \mathbb{R} -linear derivations from S to itself. Let Ω_S^1 be the S -module of regular 1-forms. Similarly define Der_F and Ω_F^1 over F . The dual inner product $I^* : V^* \times V^* \rightarrow \mathbb{R}$ naturally induces an F -bilinear form $I^* : \Omega_F^1 \times \Omega_F^1 \rightarrow F$. Then one has an F -linear bijection

$$I^* : \Omega_F^1 \rightarrow \text{Der}_F$$

^{*}Supported by JSPS Grants-in-Aid for Young Scientists (B) No. 21740014. Department of Mechanical Engineering and Science, Kyoto University, Kyoto 606-8501, Japan. email:abe.takuro.4c@kyoto-u.ac.jp

[†]Supported by JSPS Grants-in-Aid, Scientific Research (B) No. 21340001. Department of Mathematics, Hokkaido University, Sapporo, Hokkaido 060-0810, Japan. email:terao@math.sci.hokudai.ac.jp

[‡]2-8-11 Aihara, Midori-ku, Sagami-hara-shi, Kanagawa, 252-0141 Japan. email:atsushi.wakamiko@gmail.com

defined by $[I^*(\omega)](f) := I^*(\omega, df)$ for $f \in F$.

Let \mathcal{A} be an irreducible Coxeter arrangement with its Coxeter group W . For each $H \in \mathcal{A}$, choose $\alpha_H \in V^*$ with $H = \ker(\alpha_H)$. Let $Q = \prod_{H \in \mathcal{A}} \alpha_H \in S$. Recall the S -module of logarithmic forms

$$\Omega^1(\mathcal{A}, \infty) = \{\omega \in \Omega_F^1 \mid Q^N \omega \text{ and } (Q/\alpha_H)^N \omega \wedge d\alpha_H \text{ are both regular for any } H \in \mathcal{A} \text{ and } N \gg 0\}$$

and the S -module of logarithmic derivations

$$D(\mathcal{A}, -\infty) = I^*(\Omega^1(\mathcal{A}, \infty))$$

from [AT2010Z]. A map $\mathbf{m} : \mathcal{A} \rightarrow \mathbb{Z}$ is called a multiplicity. For an arbitrary multiplicity, let

$$\begin{aligned} D(\mathcal{A}, \mathbf{m}) &= \{\theta \in D(\mathcal{A}, -\infty) \mid \theta(\alpha_H) \in \alpha_H^{\mathbf{m}(H)} S_{(\alpha_H)} \text{ for all } H \in \mathcal{A}\}, \\ \Omega^1(\mathcal{A}, \mathbf{m}) &= (I^*)^{-1} D(\mathcal{A}, -\mathbf{m}), \end{aligned}$$

where $S_{(\alpha_H)}$ is the localization of S at the prime ideal (α_H) . These two modules were introduced in [Sa1980] (when \mathbf{m} is constantly equal to one), in [Z1989] (when $\text{im}(\mathbf{m}) \subset \mathbb{Z}_{>0}$), and in [A2008, AT2010Z, AT2009] (when \mathbf{m} is arbitrary). A derivation $0 \neq \theta \in \text{Der}_F$ is said to be **homogeneous of degree** d , or $\deg \theta = d$, if $\theta(\alpha) \in F$ is homogeneous of degree d whenever $\theta(\alpha) \neq 0$ ($\alpha \in V^*$). A multiarrangement $(\mathcal{A}, \mathbf{m})$ is called to be **free with exponents** $\exp(\mathcal{A}, \mathbf{m}) = (d_1, \dots, d_\ell)$ if $D(\mathcal{A}, \mathbf{m}) = \oplus_{i=1}^\ell S \cdot \theta_i$ with a homogeneous basis θ_i such that $\deg(\theta_i) = d_i$ ($i = 1, \dots, \ell$). A multiplicity $\mathbf{m} : \mathcal{A} \rightarrow \mathbb{Z}$ is said to be **equivariant** when $\mathbf{m}(H) = \mathbf{m}(wH)$ for any $H \in \mathcal{A}$ and any $w \in W$, i.e., \mathbf{m} is constant on each orbit. In this article we prove

Theorem 1.1 *For any irreducible Coxeter arrangement \mathcal{A} and any equivariant multiplicity \mathbf{m} , the multiarrangement $(\mathcal{A}, \mathbf{m})$ is free.*

For a fixed arrangement \mathcal{A} , we say that a multiplicity \mathbf{m} is **free** if $(\mathcal{A}, \mathbf{m})$ is free. Although we have a limited knowledge about the set of all free multiplicities for a fixed irreducible Coxeter arrangement \mathcal{A} , it is known that there exist infinitely many non-free multiplicities unless \mathcal{A} is either one- or two-dimensional [ATY2009]. Theorem 1.1 claims that any equivariant multiplicity is free for any irreducible Coxeter arrangement.

When the W -action on \mathcal{A} is transitive, an equivariant multiplicity is constant and a basis was constructed in [SoT1998, T2002, AY2007, AT2010Z]. So we may assume, in order to prove Theorem 1.1, that the W -action on \mathcal{A} is not transitive. In other words, we may only study the cases when \mathcal{A} is of the

type either B_ℓ, F_4, G_2 or $I_2(2n)$ ($n \geq 4$). In these cases, \mathcal{A} has exactly two W -orbits: $\mathcal{A} = \mathcal{A}_1 \cup \mathcal{A}_2$. The orbit decompositions are explicitly given by: $B_\ell = A_1^\ell \cup D_\ell$, $F_4 = D_4 \cup D_4$, $G_2 = A_2 \cup A_2$ or $I_2(2n) = I_2(n) \cup I_2(n)$ ($n \geq 4$). Note that A_1^ℓ is not irreducible.

When \mathcal{A} is irreducible, the **primitive derivations** play the central role to define the Hodge filtration introduced by K. Saito. (See [Sa2003] for example.) For $R := S^W$, let D be an element of the lowest degree in Der_R , which is called a primitive derivation corresponding to \mathcal{A} . Then D is unique up to a nonzero constant multiple. A theory of primitive derivations in the case of non-irreducible Coxeter arrangements was introduced in [AT2009]. Thus we may consider a primitive derivation D_i corresponding with the orbit \mathcal{A}_i ($1 \leq i \leq 2$). We only use D_1 because of symmetricity. Note that D_1 is not unique up to a nonzero multiple when $\mathcal{A}_1 = A_1^\ell$ (non-irreducible). Denote the reflection groups of \mathcal{A}_i by W_i ($i = 1, 2$). The Coxeter arrangements B_ℓ, F_4, G_2 and $I_2(2n)$ ($n \geq 4$) are classified into two cases, that is, (1) the primitive derivation D_1 can be chosen to be W -invariant for B_ℓ and F_4 (the first case) while (2) D_1 is W_2 -antiinvariant for G_2 and $I_2(2n)$ ($n \geq 4$) (the second case) as we will see in Section 4. Since the second cases are two-dimensional, Theorem 1.1 holds true. Thus the first case is the only remaining case to prove Theorem 1.1.

Let

$$\begin{aligned} \nabla : \text{Der}_F \times \text{Der}_F &\longrightarrow \text{Der}_F \\ (\theta, \delta) &\longmapsto \nabla_\theta \delta \end{aligned}$$

be the **Levi-Civita connection** with respect to the inner product I on V . We need the following theorem for our proof of Theorem 1.1:

Theorem 1.2 ([AT2010Z, AT2009]) *Let $D(\mathcal{A}, -\infty)^W$ be the W -invariant part of $D(\mathcal{A}, -\infty)$. Then*

$$\nabla_D : D(\mathcal{A}, -\infty)^W \xrightarrow{\sim} D(\mathcal{A}, -\infty)^W$$

is a T -linear automorphism where $T := \{f \in R \mid Df = 0\}$. When $\mathcal{A} = \mathcal{A}_1 \cup \mathcal{A}_2$ is the orbit decomposition,

$$\nabla_{D_1} : D(\mathcal{A}_1, -\infty)^{W_1} \xrightarrow{\sim} D(\mathcal{A}_1, -\infty)^{W_1}$$

is a T_1 -linear automorphism where

$$R_1 := S^{W_1}, \quad T_1 := \{f \in R_1 \mid D_1 f = 0\}.$$

Let E be the **Euler derivation** characterized by the equality $E(\alpha) = \alpha$ for every $\alpha \in V^*$. Suppose that $\mathcal{A} = \mathcal{A}_1 \cup \mathcal{A}_2$ is the orbit decomposition and that the primitive derivation D_1 is W -invariant. Define

$$E^{(p,q)} := \nabla_D^{-q} \nabla_{D_1}^{q-p} E$$

for $p, q \in \mathbb{Z}$. Here, thanks to Theorem 1.2, we may interpret $\nabla_D^m = (\nabla_D^{-1})^{-m}$ and $\nabla_{D_1}^m = (\nabla_{D_1}^{-1})^{-m}$ when m is negative. Denote the equivariant multiplicity \mathbf{m} by (m_1, m_2) when $\mathbf{m}(H) = m_1$ ($H \in \mathcal{A}_1$) and $\mathbf{m}(H) = m_2$ ($H \in \mathcal{A}_2$). Let x_1, \dots, x_ℓ be a basis for V^* and P_1, \dots, P_ℓ be a set of basic invariants with respect to W : $R = \mathbb{R}[P_1, \dots, P_\ell]$. Let $P_1^{(i)}, \dots, P_\ell^{(i)}$ be a set of basic invariants with respect to W_i : $R_i = \mathbb{R}[P_1^{(i)}, \dots, P_\ell^{(i)}]$ ($i = 1, 2$). Define

$$d_j := \deg P_j, \quad d_j^{(i)} := \deg P_j^{(i)} \quad (i = 1, 2, 1 \leq j \leq \ell).$$

We assume

$$d_1 \leq d_2 \leq \dots \leq d_\ell, \quad d_1^{(i)} \leq d_2^{(i)} \leq \dots \leq d_\ell^{(i)} \quad (i = 1, 2).$$

Then $h := d_\ell$ is called the Coxeter number of W . We call $h_i := \deg P_\ell^{(i)}$ the Coxeter number of W_i ($i = 1, 2$). We use the notation

$$\partial_{x_j} := \partial / \partial x_j, \quad \partial_{P_j} := \partial / \partial P_j, \quad \partial_{P_j^{(i)}} := \partial / \partial P_j^{(i)} \quad (1 \leq j \leq \ell, 1 \leq i \leq 2).$$

The following theorem gives an explicit construction of a basis:

Theorem 1.3 *Let \mathcal{A} be an irreducible Coxeter arrangement. Suppose that $\mathcal{A} = \mathcal{A}_1 \cup \mathcal{A}_2$ is the orbit decomposition and that the primitive derivation D_1 is W -invariant. Then*

(1) *the S -module $D(\mathcal{A}, (2p-1, 2q-1))$ is free with W -invariant basis*

$$\nabla_{\partial_{P_1}} E^{(p,q)}, \dots, \nabla_{\partial_{P_\ell}} E^{(p,q)}$$

with $\deg \nabla_{\partial_{P_i}} E^{(p,q)} = ph_1 + qh_2 - d_i + 1$ for $i = 1, \dots, \ell$,

(2) *the S -module $D(\mathcal{A}, (2p-1, 2q))$ is free with basis*

$$\nabla_{\partial_{P_1^{(1)}}} E^{(p,q)}, \dots, \nabla_{\partial_{P_\ell^{(1)}}} E^{(p,q)}$$

with $\deg \nabla_{\partial_{P_i^{(1)}}} E^{(p,q)} = ph_1 + qh_2 - d_i^{(1)} + 1$ for $i = 1, \dots, \ell$,

(3) the S -module $D(\mathcal{A}, (2p, 2q - 1))$ is free with basis

$$\nabla_{\partial_{P_1^{(2)}}} E^{(p,q)}, \dots, \nabla_{\partial_{P_\ell^{(2)}}} E^{(p,q)}$$

with $\deg \nabla_{\partial_{P_i^{(2)}}} E^{(p,q)} = ph_1 + qh_2 - d_i^{(2)} + 1$ for $i = 1, \dots, \ell$,

(4) the S -module $D(\mathcal{A}, (2p, 2q))$ is free with basis

$$\nabla_{\partial_{x_1}} E^{(p,q)}, \dots, \nabla_{\partial_{x_\ell}} E^{(p,q)}$$

with $\deg \nabla_{\partial_{x_i}} E^{(p,q)} = ph_1 + qh_2$ for $i = 1, \dots, \ell$,

The existence of the **primitive decomposition** of $D(\mathcal{A}, (2p-1, 2q-1))^W$ is proved by the following theorem:

Theorem 1.4 *Under the same assumption of Theorem 1.3 define*

$$\theta_i^{(p,q)} := \nabla_{\partial_{P_i}} E^{(p,q)} = \nabla_{\partial_{P_i}} \nabla_D^{-q} \nabla_{D_1}^{q-p} E \quad (1 \leq i \leq \ell)$$

for $p, q \in \mathbb{Z}$. Then the set

$$\{\theta_i^{(p+k, q+k)} \mid k \geq 0, 1 \leq i \leq \ell\}$$

is a T -basis for $D(\mathcal{A}, (2p-1, 2q-1))^W$. Put

$$\mathcal{G}^{(p,q)} := \bigoplus_{i=1}^{\ell} T \cdot \theta_i^{(p,q)}.$$

Then we have a T -module decomposition (called the primitive decomposition)

$$D(\mathcal{A}, (2p-1, 2q-1))^W = \bigoplus_{k \geq 0} \mathcal{G}^{(p+k, q+k)}.$$

We will also prove

Theorem 1.5 *For any irreducible Coxeter arrangement \mathcal{A} and any multiplicity \mathbf{m} , the R -module $D(\mathcal{A}, \mathbf{m})^W$ is free.*

Theorems 1.1 and 1.3 are used to prove the freeness of Shi-Catalan arrangements associated with any Weyl arrangements in [AT2010].

The organization of this article is as follows. In Section 2 we prove Theorem 1.3 when $q \geq 0$. In Section 3 we prove Theorem 1.4 to have the primitive decomposition, which is a key to complete the proof of Theorem 1.3 at the end of Section 3. In Section 4 we verify that the primitive derivation D_1 can be chosen to be W -invariant when \mathcal{A} is a Coxeter arrangement of either the type B_ℓ or F_4 . We also review the cases of G_2 and $I_2(2n)$ ($n \geq 4$) and find that the primitive derivation D_1 is W_2 -antiinvariant. In Section 5, combining Theorem 1.3 with earlier results in [T2002, AT2010Z, W2010], we finally prove Theorems 1.1 and 1.5.

2 Proof of Theorem 1.3 when $q \geq 0$

In this section we prove Theorem 1.3 when $q \geq 0$.

Recall $R = S^W = \mathbb{R}[P_1, \dots, P_\ell]$ is the invariant ring with basic invariants P_1, \dots, P_ℓ such that $2 = \deg P_1 < \deg P_2 \leq \dots \leq \deg P_{\ell-1} < \deg P_\ell = h$, where h is the Coxeter number of W . Put $D = \partial_{P_\ell} \in \text{Der } R$ which is a primitive derivation. Recall $T = \ker(D : R \rightarrow R) = \mathbb{R}[P_1, \dots, P_{\ell-1}]$. Then the covariant derivative ∇_D is T -linear. For $\mathbf{P} := [P_1, \dots, P_\ell]$, the Jacobian matrix $J(\mathbf{P})$ is defined as the matrix whose (i, j) -entry is $\frac{\partial P_j}{\partial x_i}$. Define $A := [I^*(dx_i, dx_j)]_{1 \leq i, j \leq \ell}$ and $G := [I^*(dP_i, dP_j)]_{1 \leq i, j \leq \ell} = J(\mathbf{P})^T A J(\mathbf{P})$.

Definition 2.1 ([Y2002, W2010]) Let $\mathbf{m} : \mathcal{A} \rightarrow \mathbb{Z}$ and $\zeta \in D(\mathcal{A}, -\infty)^W$. We say that ζ is **m-universal** when ζ is homogeneous and the S -linear map

$$\begin{aligned} \Psi_\zeta : \text{Der}_S &\longrightarrow D(\mathcal{A}, 2\mathbf{m}) \\ \theta &\longmapsto \nabla_\theta \zeta \end{aligned}$$

is bijective.

Example 2.2 The Euler derivation E is **0-universal** because $\Psi_E(\delta) = \nabla_\delta E = \delta$ and $D(\mathcal{A}, \mathbf{0}) = \text{Der}_S$.

Recall the T -automorphisms

$$\nabla_D^k : D(\mathcal{A}, -\infty)^W \xrightarrow{\sim} D(\mathcal{A}, -\infty)^W \quad (k \in \mathbb{Z})$$

from Theorem 1.2. Recall the following two results concerning the **m-universality**:

Theorem 2.3 ([W2010, Theorem 2.8]) If ζ is **m-universal**, then $\nabla_D^{-1}\zeta$ is **(m + 1)-universal**.

Proposition 2.4 ([W2010, Proposition 2.7]) Suppose that ζ is **m-universal**. Let $\mathbf{k} : \mathcal{A} \rightarrow \{+1, 0, -1\}$. Then an S -homomorphism

$$\Phi_\zeta : D(\mathcal{A}, \mathbf{k}) \rightarrow D(\mathcal{A}, \mathbf{k} + 2\mathbf{m})$$

defined by

$$\Phi_\zeta(\theta) := \nabla_\theta \zeta$$

gives an S -module isomorphism.

We require that assumption of Theorem 1.3 is satisfied in the rest of this section: Suppose that $\mathcal{A} = \mathcal{A}_1 \cup \mathcal{A}_2$ is the orbit decomposition and that D_1 , a primitive derivation with respect to \mathcal{A}_1 in the sense of [AT2009, Definition 2.4], is W -invariant. Let W_i , R_i , $P_j^{(i)}$, T_i , D_i ($i = 1, 2$) are defined as in Section 1. Even when \mathcal{A}_1 is not irreducible, we may consider a T_1 -isomorphism

$$\nabla_{D_1}^k : D(\mathcal{A}_1, -\infty)^{W_1} \xrightarrow{\sim} D(\mathcal{A}_1, -\infty)^{W_1} \quad (k \in \mathbb{Z})$$

from Theorem 1.2.

Proposition 2.5 *Suppose $q \geq 0$. The derivation $E^{(p,q)} := \nabla_D^{-q} \nabla_{D_1}^{q-p} E$ is (p, q) -universal.*

Proof. When \mathcal{A}_1 is irreducible, [AY2007] and [AT2010Z] imply that $\nabla_{D_1}^{q-p} E$ is $(p - q, 0)$ -universal. When \mathcal{A}_1 is not irreducible, $\nabla_{D_1}^{q-p} E$ is $(p - q, 0)$ -universal because of [AT2009]. Thus $E^{(p,q)} = \nabla_D^{-q} \nabla_{D_1}^{q-p} E$ is (p, q) -universal by Theorem 2.3. \square

Since $E^{(p,q)}$ is (p, q) -universal, Proposition 2.4 yields the following:

Proposition 2.6 *Let $q \geq 0$ and $\mathbf{m} : \mathcal{A} \rightarrow \{+1, 0, -1\}$. Then an S -homomorphism*

$$\Phi_{p,q} : D(\mathcal{A}, \mathbf{m}) \rightarrow D(\mathcal{A}, (2p, 2q) + \mathbf{m})$$

defined by

$$\Phi_{p,q}(\theta) := \nabla_\theta E^{(p,q)}$$

gives an S -module isomorphism.

Proof of Theorem 1.3 ($q \geq 0$). We may apply Proposition 2.6 because

- (1) $\partial_{P_1}, \dots, \partial_{P_\ell}$ form a basis for $D(\mathcal{A}, (-1, -1))$,
- (2) $\partial_{P_1^{(1)}}, \dots, \partial_{P_\ell^{(1)}}$ form a basis for $D(\mathcal{A}, (-1, 0))$,
- (3) $\partial_{P_1^{(2)}}, \dots, \partial_{P_\ell^{(2)}}$ form a basis for $D(\mathcal{A}, (0, -1))$, and
- (4) $\partial_{x_1}, \dots, \partial_{x_\ell}$ form a basis for $D(\mathcal{A}, (0, 0))$. \square

3 Primitive decompositions

In this section we first prove Theorem 1.4 to define the primitive decomposition of $D(\mathcal{A}, (2p - 1, 2q - 1))^W$. Next we prove Theorem 1.3.

Proposition 3.1 *Let ζ be \mathbf{m} -universal. Then*

- (1) *the set $\{\nabla_{\partial_{P_j}} \nabla_D^{-k} \zeta \mid 1 \leq j \leq \ell, k \geq 0\}$ is linearly independent over T .*
- (2) *Define $\mathcal{G}^{(k)}$ to be the free T -module with basis $\{\nabla_{\partial_{P_j}} \nabla_D^{-k} \zeta \mid 1 \leq j \leq \ell\}$ for $k \geq 0$. Then the Poincaré series $\text{Poin}(\bigoplus_{k \geq 0} \mathcal{G}^{(k)}, t)$ satisfies:*

$$\text{Poin}(\bigoplus_{k \geq 0} \mathcal{G}^{(k)}, t) = \left(\prod_{i=1}^{\ell} \frac{1}{1 - t^{d_i}} \right) \left(\sum_{j=1}^{\ell} t^{p-d_j} \right),$$

where $p = \deg \zeta$ and $d_j = \deg P_j$ ($1 \leq j \leq \ell$).

(3)

$$D(\mathcal{A}, 2\mathbf{m} - 1)^W = \bigoplus_{k \geq 0} \mathcal{G}^{(k)}.$$

Proof. Let $k \in \mathbb{Z}_{\geq 0}$. By Theorem 2.3, $\zeta^{(k)} := \nabla_D^{-k} \zeta$ is $(\mathbf{m} + k)$ -universal, where the “ k ” in the $(\mathbf{m} + k)$ stands for the constant multiplicity k by abuse of notation. Thus by Proposition 2.4 we have the following two bases:

$$\nabla_{\partial_{P_1}} \zeta^{(k)}, \dots, \nabla_{\partial_{P_\ell}} \zeta^{(k)},$$

for the S -module $D(\mathcal{A}, 2\mathbf{m} + 2k - 1)$ and

$$\nabla_{\partial_{I^*(dP_1)}} \zeta^{(k)}, \dots, \nabla_{\partial_{I^*(dP_\ell)}} \zeta^{(k)},$$

for the S -module $D(\mathcal{A}, 2\mathbf{m} + 2k + 1)$. Note that the two bases are also R -bases for $D(\mathcal{A}, 2\mathbf{m} + 2k - 1)^W$ and $D(\mathcal{A}, 2\mathbf{m} + 2k + 1)^W$ respectively. Since the T -automorphism

$$\nabla_D: D(\mathcal{A}, -\infty)^W \xrightarrow{\sim} D(\mathcal{A}, -\infty)^W$$

in Theorem 1.2 induces a T -linear bijection

$$\nabla_D: D(\mathcal{A}, 2\mathbf{m} + 2k + 1)^W \xrightarrow{\sim} D(\mathcal{A}, 2\mathbf{m} + 2k - 1)^W$$

as in [AT2009, Theorem 4.4], we may find an $\ell \times \ell$ -matrix $B^{(k)}$ with entries in R such that

$$\begin{aligned} \nabla_D \left(\left[\nabla_{\partial_{P_1}} \zeta^{(k)}, \dots, \nabla_{\partial_{P_\ell}} \zeta^{(k)} \right] G \right) &= \nabla_D \left[\nabla_{\partial_{I^*(dP_1)}} \zeta^{(k)}, \dots, \nabla_{\partial_{I^*(dP_\ell)}} \zeta^{(k)} \right] \\ &= \left[\nabla_{\partial_{P_1}} \zeta^{(k)}, \dots, \nabla_{\partial_{P_\ell}} \zeta^{(k)} \right] B^{(k)}. \end{aligned}$$

The degree of (i, j) -th entry of $B^{(k)}$ is $m_i + m_j - h \leq h - 2 < h$. In particular, the degree of $B_{i, \ell+1-i}^{(k)}$ is 0 and $B_{i, j}^{(k)} = 0$ if $i + j < \ell + 1$. Hence each entry

of $B^{(k)}$ lies in T and $\det B^{(k)} \in \mathbb{R}$. Since D is a derivation of the minimum degree in Der_R , one gets $[D, \partial_{P_i}] = 0$. Thus $\nabla_D \nabla_{\partial_{P_i}} = \nabla_{\partial_{P_i}} \nabla_D$. Operate ∇_D^{-1} on the both sides of the equality above, and get

$$\left[\nabla_{\partial_{P_1}} \zeta^{(k)}, \dots, \nabla_{\partial_{P_\ell}} \zeta^{(k)} \right] G = \left[\nabla_{\partial_{P_1}} \zeta^{(k+1)}, \dots, \nabla_{\partial_{P_\ell}} \zeta^{(k+1)} \right] B^{(k)}.$$

This implies that $\det B^{(k)} \in \mathbb{R}^\times$ because $\nabla_{\partial_{P_1}} \zeta^{(k)}, \dots, \nabla_{\partial_{P_\ell}} \zeta^{(k)}$ are linearly independent over S . Inductively we have

$$\begin{aligned} \left[\nabla_{\partial_{P_1}} \zeta^{(k+1)}, \dots, \nabla_{\partial_{P_\ell}} \zeta^{(k+1)} \right] &= \left[\nabla_{\partial_{P_1}} \zeta^{(k)}, \dots, \nabla_{\partial_{P_\ell}} \zeta^{(k)} \right] G(B^{(k)})^{-1} \\ &= \left[\nabla_{\partial_{P_1}} \zeta, \dots, \nabla_{\partial_{P_\ell}} \zeta \right] G(B^{(0)})^{-1} G(B^{(1)})^{-1} \dots G(B^{(k)})^{-1} \\ &= \left[\nabla_{\partial_{P_1}} \zeta, \dots, \nabla_{\partial_{P_\ell}} \zeta \right] G_{k+1}, \end{aligned}$$

where $G_i = G(B^{(0)})^{-1} G(B^{(1)})^{-1} \dots G(B^{(i-1)})^{-1}$ ($i \geq 0$). Note that G appears i times in the definition of G_i . For $M = (m_{ij}) \in M_\ell(F)$, define $D[M] = (D(m_{ij})) \in M_\ell(F)$. Then $D^j[G_i] = 0$ when $j > i$ and $\det D^i[G_i] \neq 0$ because $\det D[G] \neq 0$ and $D^2[G] = 0$ (e.g., see [Sa1993, AT2009]).

(1) Suppose that $\{\nabla_{\partial_{P_j}} \zeta^{(k)} \mid 1 \leq j \leq \ell, k \geq 0\}$ is linearly dependent over T . Then there exist ℓ -dimensional column vectors $\mathbf{g}_0, \mathbf{g}_1, \dots, \mathbf{g}_q \in T^\ell$ ($q \geq 0$) with $\mathbf{g}_q \neq \mathbf{0}$ such that

$$\mathbf{0} = \sum_{i=0}^q \left[\nabla_{\partial_{P_1}} \zeta^{(i)}, \dots, \nabla_{\partial_{P_\ell}} \zeta^{(i)} \right] \mathbf{g}_i = \left[\nabla_{\partial_{P_1}} \zeta, \dots, \nabla_{\partial_{P_\ell}} \zeta \right] \left(\sum_{i=0}^q G_i \mathbf{g}_i \right).$$

Since $\nabla_{\partial_{P_1}} \zeta, \dots, \nabla_{\partial_{P_\ell}} \zeta$ are linearly independent over R , one has

$$\mathbf{0} = \sum_{i=0}^q G_i \mathbf{g}_i.$$

Applying the operator D on the both sides q times, we get $D^q[G_q] \mathbf{g}_q = \mathbf{0}$. Thus $\mathbf{g}_q = \mathbf{0}$ which is a contradiction. This proves (1).

(2) Compute

$$\begin{aligned} \text{Poin}\left(\bigoplus_{k \geq 0} \mathcal{G}^{(k)}, t\right) &= \sum_{k \geq 0} \left(\prod_{i=1}^{\ell-1} \frac{1}{1-t^{d_i}} \right) \left(\sum_{j=1}^{\ell} t^{p-d_j+kd_\ell} \right) \\ &= \left(\prod_{i=1}^{\ell-1} \frac{1}{1-t^{d_i}} \right) \left(\sum_{k \geq 0} t^{kd_\ell} \right) \left(\sum_{j=1}^{\ell} t^{p-d_j} \right) \\ &= \left(\prod_{i=1}^{\ell} \frac{1}{1-t^{d_i}} \right) \left(\sum_{j=1}^{\ell} t^{p-d_j} \right). \end{aligned}$$

(3) We have

$$D(\mathcal{A}, 2\mathbf{m} - 1)^W \supseteq \bigoplus_{k \geq 0} \mathcal{G}^{(k)}$$

by (1). So it suffices to prove

$$\text{Poin}(D(\mathcal{A}, 2\mathbf{m} - 1)^W, t) = \text{Poin}\left(\bigoplus_{k \geq 0} \mathcal{G}^{(k)}, t\right).$$

Since $D(\mathcal{A}, 2\mathbf{m} - 1)^W$ is a free R -module with a basis

$$\nabla_{\partial_{P_1}} \zeta, \dots, \nabla_{\partial_{P_\ell}} \zeta,$$

we obtain

$$\text{Poin}(D(\mathcal{A}, 2\mathbf{m} - 1)^W, t) = \left(\prod_{i=1}^{\ell} \frac{1}{1 - t^{d_i}} \right) \left(\sum_{i=1}^{\ell} t^{p-d_j} \right) = \text{Poin}\left(\bigoplus_{k \geq 0} \mathcal{G}^{(k)}, t\right),$$

which completes the proof. \square

We require that the assumption of Theorem 1.3 is satisfied in the rest of this section.

Proof of Theorem 1.4. Suppose $q \geq 0$ to begin with. Then, by Proposition 3.4, $E^{(p,q)}$ is (p, q) -universal. Apply Proposition 3.1 for $\zeta = E^{(p,q)}$ and $\mathbf{m} = (p, q)$, and we have Theorem 1.4:

$$D(\mathcal{A}, (2p - 1, 2q - 1))^W = \bigoplus_{k \geq 0} \mathcal{G}^{(p+k, q+k)}$$

when $q \geq 0$. Send the both handsides by ∇_D , and we get

$$D(\mathcal{A}, (2p - 3, 2q - 3))^W = \bigoplus_{k \geq 0} \mathcal{G}^{(p+k-1, q+k-1)}$$

because $\nabla_D (D(\mathcal{A}, (2p - 1, 2q - 1))^W) = D(\mathcal{A}, (2p - 3, 2q - 3))^W$ as in [AT2009, Theorem 4.4] and $\nabla_D(\theta_i^{(p,q)}) = \theta_i^{(p-1, q-1)}$. Apply ∇_D repeatedly to complete the proof for all $q \in \mathbb{Z}$. \square

Note that we do not assume $p \geq 0$ in the following proposition:

Proposition 3.2 *For $p, q \in \mathbb{Z}$, the S -module $D(\mathcal{A}, (2p - 1, 2q - 1))$ has a W -invariant basis.*

Proof. Recall that

$$\nabla_{\partial_{P_1}} E^{(p,q)}, \nabla_{\partial_{P_2}} E^{(p,q)}, \dots, \nabla_{\partial_{P_\ell}} E^{(p,q)},$$

which are W -invariant, form an S -basis for $D(\mathcal{A}, (2p-1, 2q-1))$ when $q \geq 0$ by Theorem 1.3 (1). It is then easy to see that they are also an R -basis for $D(\mathcal{A}, (2p-1, 2q-1))^W$ for $q \geq 0$. By [A2008] [AT2010Z], there exists a W -equivariant nondegenerate S -bilinear pairing

$$(\ , \) : D(\mathcal{A}, (2p-1, 2q-1)) \times D(\mathcal{A}, (-2p+1, -2q+1)) \longrightarrow S,$$

characterized by

$$(I^*(\omega), \theta) = \langle \omega, \theta \rangle$$

where $\omega \in \Omega^1(\mathcal{A}, (-2p+1, -2q+1))$ and $\theta \in D(\mathcal{A}, (-2p+1, -2q+1))$. Let $\theta_1, \dots, \theta_\ell$ denote the dual basis for $D(\mathcal{A}, (-2p+1, -2q+1))$ satisfying

$$\left(\nabla_{\partial_{P_i}} E^{(p,q)}, \theta_j \right) = \delta_{ij}$$

for $1 \leq i, j \leq \ell$. Then $\theta_1, \dots, \theta_\ell$ are W -invariant because the pairing $(\ , \)$ is W -equivariant. \square

Although the following lemma is standard and easy, we give a proof for completeness.

Lemma 3.3 *Let M be an S -submodule of Der_F . The following two conditions are equivalent:*

- (1) *M has a W -invariant basis Θ over S .*
- (2) *The W -invariant part M^W is a free R -module with a basis Θ and there exists a natural S -linear isomorphism*

$$M^W \otimes_R S \simeq M.$$

Proof. It suffices to prove that (1) implies (2) because the other implication is obvious. Suppose that $\Theta = \{\theta_\lambda\}_{\lambda \in \Lambda}$ is a W -invariant basis for M over S . Since it is linearly independent over S , so is over R . Let $\theta \in M^W$. Express

$$\theta = \sum_{i=1}^n f_i \theta_i$$

with $f_i \in S$ and $\theta_i \in \Theta$ ($i = 1, \dots, n$). Let $w \in W$ act on the both handsides. Then we get

$$\theta = \sum_{i=1}^n w(f_i) \theta_i.$$

This implies $f_i = w(f_i)$ for every $w \in W$. Hence $f_i \in R$ for each i . Therefore Θ is a basis for M^W over R . This is (2). \square

Proposition 3.4 *For any $p, q \in \mathbb{Z}$, $E^{(p,q)}$ is (p, q) -universal.*

Proof. By Theorem 1.4 we have the decomposition:

$$D(\mathcal{A}, (2p-1, 2q-1))^W = \bigoplus_{k \geq 0} \mathcal{G}^{(p+k, q+k)}$$

for $p, q \in \mathbb{Z}$. As we saw in Proposition 3.1 (2), we have

$$(3.1) \quad \text{Poin}(D(\mathcal{A}, (2p-1, 2q-1))^W, t) = \text{Poin}\left(\bigoplus_{k \geq 0} \mathcal{G}^{(p+k, q+k)}, t\right) \\ = \left(\prod_{i=1}^{\ell} \frac{1}{1-t^{d_i}}\right) \left(\sum_{i=1}^{\ell} t^{m-d_j}\right),$$

where $m := \deg E^{(p,q)}$. Recall that the S -module $D(\mathcal{A}, (2p-1, 2q-1))$ has a W -invariant basis $\theta_1, \dots, \theta_{\ell}$ by Proposition 3.2. By Lemma 3.3, we know that $\theta_1, \dots, \theta_{\ell}$ form a basis for the R -module $D(\mathcal{A}, (2p-1, 2q-1))^W$. Thanks to (3.1) we may assume that $\deg \theta_j = m - d_j = \deg \nabla_{\partial_{P_j}} E^{(p,q)}$. Therefore there exists $M \in M_{\ell}(R)$ such that

$$[\theta_1, \dots, \theta_{\ell}]M = [\nabla_{\partial_{P_1}} E^{(p,q)}, \dots, \nabla_{\partial_{P_{\ell}}} E^{(p,q)}]$$

with $\det M \in \mathbb{R}$. Since

$$\max_{1 \leq i, j \leq \ell} \left| \deg \theta_i - \deg \nabla_{\partial_{P_j}} E^{(p,q)} \right| = d_{\ell} - d_1 < \deg P_{\ell},$$

we get $M \in M_{\ell}(T)$. Since $\nabla_{\partial_{P_1}} E^{(p,q)}, \dots, \nabla_{\partial_{P_{\ell}}} E^{(p,q)}$ are linearly independent over T by Proposition 3.1 (1), we have $\det M \in \mathbb{R}^{\times}$. Thus

$$\nabla_{\partial_{P_1}} E^{(p,q)}, \dots, \nabla_{\partial_{P_{\ell}}} E^{(p,q)}$$

form an S -basis for $D(\mathcal{A}, (2p-1, 2q-1))$. Since

$$\left[\nabla_{\partial_{P_1}} E^{(p,q)}, \dots, \nabla_{\partial_{P_{\ell}}} E^{(p,q)} \right] J(\mathbf{P})^T = \left[\nabla_{\partial_{x_1}} E^{(p,q)}, \dots, \nabla_{\partial_{x_{\ell}}} E^{(p,q)} \right],$$

we may apply the multi-arrangement version of Saito's criterion [Sa1980, Z1989, A2008] to prove that $\nabla_{\partial_{x_1}} E^{(p,q)}, \dots, \nabla_{\partial_{x_{\ell}}} E^{(p,q)}$ form an S -basis for $D(\mathcal{A}, (2p, 2q))$ for any $p, q \in \mathbb{Z}$. This shows that $E^{(p,q)}$ is (p, q) -universal for any $p, q \in \mathbb{Z}$. \square

Proof of Theorem 1.3 ($q \in \mathbb{Z}$). Theorem 2.3 and Proposition 3.4 complete the proof by the same argument as that in Section 2 for $q \geq 0$. \square

4 The cases of B_ℓ , F_4 , G_2 and $I_2(2n)$

• The case of B_ℓ

The roots of the type B_ℓ are:

$$\pm x_i, \pm x_i \pm x_j \quad (1 \leq i < j \leq \ell)$$

in terms of an orthonormal basis x_1, \dots, x_ℓ for V^* . Altogether there are $2\ell^2$ of them. Define

$$Q_1 := \prod_{i=1}^{\ell} x_i, \quad Q_2 := \prod_{1 \leq i < j \leq \ell} (x_i \pm x_j), \quad Q = Q_1 Q_2.$$

Then the arrangement \mathcal{A}_1 defined by Q_1 is of the type $A_1 \times \dots \times A_1 = A_1^\ell$. The arrangement \mathcal{A}_2 defined by Q_2 is of the type D_ℓ . The arrangement \mathcal{A} defined by Q is of the type B_ℓ and $\mathcal{A} = \mathcal{A}_1 \cup \mathcal{A}_2$ is the orbit decomposition. Note that A_1^ℓ is not irreducible. Define

$$D_1 := \sum_{i=1}^{\ell} \frac{1}{x_i} \partial_{x_i}$$

which is a primitive derivation in the sense of [AT2009]. Obviously D_1 is W -invariant. Let $P_j = \sum_{i=1}^{\ell} x_i^{2j}$ ($j \geq 1$). Then P_1, \dots, P_ℓ form a set of basic invariants under W while $Q_1, P_1, \dots, P_{\ell-1}$ form a set of basic invariants under W_2 . Define a primitive derivation D_2 with respect to \mathcal{A}_2 so that

$$D_2(Q_1) = D_2(P_j) = 0 \quad (j = 1, \dots, \ell - 2), \quad D_2(P_{\ell-1}) = 1.$$

Thus

$$(wD_2)(P_{\ell-1}) = D_2(w^{-1}P_{\ell-1}) = D_2(P_{\ell-1}) = 1 \quad (w \in W).$$

This implies that D_2 is W -invariant.

• The case of F_4

The roots of the type F_4 are:

$$\pm x_i, (\pm x_1 \pm x_2 \pm x_3 \pm x_4)/2, \pm x_i \pm x_j \quad (1 \leq i < j \leq 4)$$

in terms of an orthonormal basis x_1, x_2, x_3, x_4 for V^* . Altogether there are 48 of them. Define

$$Q_1 := \prod_{1 \leq i < j \leq 4} (x_i \pm x_j), \quad Q_2 := \prod_{i=1}^4 x_i \prod (x_1 \pm x_2 \pm x_3 \pm x_4), \quad Q = Q_1 Q_2.$$

The arrangement \mathcal{A}_i defined by Q_i is of the type D_4 ($i = 1, 2$). Then the arrangement \mathcal{A} defined by Q is of the type F_4 and $\mathcal{A} = \mathcal{A}_1 \cup \mathcal{A}_2$ is the orbit decomposition. Define

$$P_1^{(1)} = \sum_{i=1}^4 x_i^2, \quad P_2^{(1)} = \sum_{i=1}^4 x_i^4, \quad P_3^{(1)} = x_1 x_2 x_3 x_4, \quad P_4^{(1)} = \sum_{i=1}^4 x_i^6 + 5 \sum_{i \neq j} x_i^2 x_j^4.$$

Compute

$$P_4^{(1)} = -4 \sum_{i=1}^4 x_i^6 + 5 P_1^{(1)} P_2^{(1)}.$$

Thus $P_1^{(1)}, P_2^{(1)}, P_3^{(1)}, P_4^{(1)}$ are a set of basic invariants under W_1 . The reflection τ with respect to $x_1 + x_2 + x_3 + x_4 = 0$ is given by

$$\tau(x_i) = \frac{2x_i - \sum_{j=1}^4 x_j}{2} \quad (i = 1, 2, 3, 4).$$

A calculation shows that $P_4^{(1)}$ is τ -invariant. Let s_i denote the reflection with respect to $x_i = 0$ ($1 \leq i \leq 4$). Since the Coxeter group W_2 is generated by τ and s_i ($1 \leq i \leq 4$), we know that $P_4^{(1)}$ is W_2 -invariant thus W -invariant. Define a primitive derivation D_1 with respect to \mathcal{A}_1 so that

$$D_1(P_j^{(1)}) = 0 \quad (j = 1, 2, 3), \quad D_1(P_4^{(1)}) = 1.$$

Thus

$$(wD_1)(P_4^{(1)}) = D_1(w^{-1}P_4^{(1)}) = D_1(P_4^{(1)}) = 1 \quad (w \in W).$$

This implies that D_1 is W -invariant. We conclude that D_2 is also W -invariant because an orthonormal coordinate change

$$x_1 = \frac{y_1 - y_2}{\sqrt{2}}, \quad x_2 = \frac{y_1 + y_2}{\sqrt{2}}, \quad x_3 = \frac{y_3 - y_4}{\sqrt{2}}, \quad x_4 = \frac{y_3 + y_4}{\sqrt{2}}$$

switches \mathcal{A}_1 and \mathcal{A}_2 .

• **The cases of G_2 and $I_2(2n)$ ($n \geq 4$)**

The arrangement \mathcal{A} of the type G_2 has exactly two orbits \mathcal{A}_1 and \mathcal{A}_2 , each of which is of the type A_2 . Let $n \geq 4$. Then the arrangement \mathcal{A} of the type $I_2(2n)$ has exactly two orbits \mathcal{A}_1 and \mathcal{A}_2 , each of which is of the type $I_2(n)$. In both cases, by [W2010], one may choose

$$D_1 = Q_2 D, \quad D_2 = Q_1 D.$$

Since Q_2 is W_2 -antiinvariant and D is W -invariant, D_1 is W_2 -antiinvariant. Similarly D_2 is W_1 -antiinvariant.

5 Proofs of Theorems 1.1 and 1.5

Assume that \mathcal{A} is an irreducible Coxeter arrangement in the rest of the article.

Proof of Theorem 1.1. If \mathcal{A} has the single orbit, then the result in [T2002, AY2007, AT2010Z] completes the proof. If not, then \mathcal{A} has exactly two orbits. If \mathcal{A} is of the type either G_2 or $I_2(2n)$ with $n \geq 4$, then $D(\mathcal{A}, \mathbf{m})$ is a free S -module because \mathcal{A} lies in a two-dimensional vector space. For the remaining cases of the type B_ℓ and F_4 , Section 4 allows us to apply Theorem 1.3 to complete the proof. \square

A multiplicity $\mathbf{m} : \mathcal{A} \rightarrow \mathbb{Z}$ is said to be **odd** if its image lies in $1 + 2\mathbb{Z}$.

Proposition 5.1 *If \mathbf{m} is equivariant and odd, then $D(\mathcal{A}, \mathbf{m})$ has a W -invariant basis over S .*

Proof. When \mathcal{A} has the single orbit, \mathbf{m} is constant. In this case Proposition was proved in [T2002, AY2007, AT2010Z]. If \mathcal{A} is of the type either G_2 or $I_2(2n)$ ($n \geq 4$), then Proposition was verified in [W2010]. For the remaining cases of B_ℓ and F_4 , Proposition 3.2 completes the proof. \square

Recall the W -action on \mathcal{A} :

$$W \times \mathcal{A} \longrightarrow \mathcal{A}$$

by sending (w, H) to wH ($w \in W$, $H \in \mathcal{A}$). For any multiplicity $\mathbf{m} : \mathcal{A} \rightarrow \mathbb{Z}$, define a new multiplicity \mathbf{m}^* by

$$\mathbf{m}^*(H) := \max_{w \in W} (2 \cdot \lfloor \mathbf{m}(wH)/2 \rfloor + 1),$$

where $\lfloor a \rfloor$ stands for the greatest integer not exceeding a . Then \mathbf{m}^* is obviously equivariant and odd.

Proposition 5.2 *For any irreducible Coxeter arrangement \mathcal{A} and any multiplicity \mathbf{m} ,*

$$D(\mathcal{A}, \mathbf{m})^W = D(\mathcal{A}, \mathbf{m}^*)^W.$$

Proof. Since $\mathbf{m}(H) \leq \mathbf{m}^*(H)$ for any $H \in \mathcal{A}$, we have

$$D(\mathcal{A}, \mathbf{m})^W \supseteq D(\mathcal{A}, \mathbf{m}^*)^W.$$

We will show the other inclusion. Let $H \in \mathcal{A}$ and $\theta \in D(\mathcal{A}, \mathbf{m})^W$. It suffices to verify the following two statements:

$$(A) \quad \theta(\alpha_H) \in \alpha_H^{\mathbf{m}(wH)} S_{(\alpha_H)} \text{ for any } w \in W,$$

(B) $\theta(\alpha_H) \in \alpha_H^{2m} S_{(\alpha_H)}$ implies $\theta(\alpha_H) \in \alpha_H^{2m+1} S_{(\alpha_H)}$ for any $m \in \mathbb{Z}$.

For $w \in W$ let w^{-1} act on the both sides of

$$\theta(\alpha_{wH}) \in \alpha_{wH}^{\mathbf{m}(wH)} S_{(\alpha_{wH})}$$

to get

$$\theta(\alpha_H) \in \alpha_H^{\mathbf{m}(wH)} S_{(\alpha_H)}.$$

This verifies (A).

Fix $H \in \mathcal{A}$. Let s be the orthogonal reflection through H . Then $s(\alpha_H) = -\alpha_H$. Suppose that $\theta(\alpha_H) = \alpha_H^{2m} p$ with $p \in S_{(\alpha_H)}$. Let s act on the both handsides and we have $\theta(-\alpha_H) = (-\alpha_H)^{2m} s(p)$. This implies $-p = s(p)$. Since $s(p) = p$ on H , one has $p = 0$ on H , which implies $p \in \alpha_H S_{(\alpha_H)}$. This verifies (B). \square

Proof of Theorem 1.5. Thanks to Proposition 5.2 we may assume that \mathbf{m} is equivariant and odd. Apply Proposition 5.1 and Lemma 3.3. \square

Corollary 5.3

$$D(\mathcal{A}, \mathbf{m})^W \otimes_R S \simeq D(\mathcal{A}, \mathbf{m}^*).$$

Proof. Apply Proposition 5.1 and Lemma 3.3 to get

$$D(\mathcal{A}, \mathbf{m}^*)^W \otimes_R S \simeq D(\mathcal{A}, \mathbf{m}^*).$$

Then Proposition 5.2 completes the proof. \square

The following corollary shows that the converse of Proposition 5.1 is true.

Corollary 5.4 *The S -module $D(\mathcal{A}, \mathbf{m})$ has a W -invariant basis if and only if \mathbf{m} is odd and equivariant.*

Proof. Assume that $D(\mathcal{A}, \mathbf{m})$ has a W -invariant basis over S . Then, by Lemma 3.3, we get

$$D(\mathcal{A}, \mathbf{m})^W \otimes_R S \simeq D(\mathcal{A}, \mathbf{m}).$$

Compare this with Corollary 5.3 and we know that there exists a common S -basis for both $D(\mathcal{A}, \mathbf{m})$ and $D(\mathcal{A}, \mathbf{m}^*)$. By the multi-arrangement version of Saito's criterion [Sa1980, Z1989, A2008], we have $\mathbf{m} = \mathbf{m}^*$. \square

References

- [A2008] T. Abe, A generalized logarithmic module and duality of Coxeter multiarrangements. arXiv.0807.2552v1.
- [AT2010Z] T. Abe and H. Terao, A primitive derivation and logarithmic differential forms of Coxeter arrangements. *Math. Z.* **264** (2010), 813–828.
- [AT2009] T. Abe and H. Terao, Primitive filtrations of the modules of invariant logarithmic forms of Coxeter arrangements. arXiv:0910.2506v1, to appear in *J. Algebra*.
- [AT2010] T. Abe and H. Terao, The freeness of Shi-Catalan arrangements. In preparation.
- [ATY2009] T. Abe, H. Terao and M. Yoshinaga, Totally free arrangements of hyperplanes. *Proc. AMS* **137** (2009), 1405–1410.
- [AY2007] T. Abe and M. Yoshinaga, Coxeter multiarrangements with quasi-constant multiplicities. *J. Algebra* **322** (2009), 2839–2847.
- [OT1992] P. Orlik and H. Terao, *Arrangements of hyperplanes*. Grundlehren der Mathematischen Wissenschaften, **300**. Springer-Verlag, Berlin, 1992.
- [Sa1980] K. Saito, Theory of logarithmic differential forms and logarithmic vector fields. *J. Fac. Sci. Univ. Tokyo Sect. IA Math.* **27** (1980), 265–291.
- [Sa1993] K. Saito, On a linear structure of the quotient variety by a finite reflection group. *Publ. RIMS, Kyoto Univ.* **29** (1993), 535–579.
- [Sa2003] K. Saito, Uniformization of the orbifold of a finite reflection group. *RIMS preprint* **1414** (2003).
- [SoT1998] L. Solomon and H. Terao, The double Coxeter arrangements. *Comment. Math. Helv.* **73** (1998), 237–258.
- [T2002] H. Terao, Multiderivations of Coxeter arrangements. *Invent. Math.* **148** (2002), 659–674.
- [W2010] A. Wakamiko, Bases for the derivation modules of two-dimensional multi-Coxeter arrangements and universal derivations. arXiv.1010.5266v1, to appear in *Hokkaido Math. J.*

- [Y2002] M. Yoshinaga, The primitive derivation and freeness of multi-Coxeter arrangements. *Proc. Japan Acad. Ser. A* **78** (2002), no. 7, 116–119.
- [Z1989] G. Ziegler, Multiarrangements of hyperplanes and their freeness. *Singularities* (Iowa City, IA, 1986), 345–359, *Contemp. Math.*, **90**, Amer. Math. Soc., Providence, RI, 1989.